arXiv:1107.1368v2 [cond-mat.stat-mech] 16 Sep 2011

Correlations in the Ising antiferromagnet on the anisotropic kagome lattice

Walter Apel[1] **and Hans-Ulrich Everts**[2]

[1] Physikalisch-Technische Bundesanstalt (PTB), Bundesallee 100, 38116
Braunschweig, Germany
[2] Institut für Theoretische Physik, Leibniz Universität Hannover, Appelstraße 2,
30167 Hannover, Germany

E-mail: `Walter.Apel@ptb.de` and `everts@itp.uni-hannover.de`

Abstract. We study the correlation function of middle spins, i. e. of spins on intermediate sites between two adjacent parallel lattice axes, of the spatially anisotropic Ising antiferromagnet on the kagome lattice. It is given rigorously by a Toeplitz determinant. The large-distance behaviour of this correlation function is obtained by analytic methods. For shorter distances we evaluate the Toeplitz determinant numerically. The correlation function is found to vanish exactly on a line $J_d(T)$ in the $T - J$ (temperature vs. coupling constant) phase diagram. This disorder line divides the phase diagram into two regions. For $J < J_d(T)$ the correlations display the features of an unfrustrated two-dimensional Ising magnet, whereas for $J > J_d(T)$ the correlations between the middle spins are seen to be strongly influenced by the short-range antiferromagnetic order that prevails among the spins of the adjacent lattice axes. While for $J < J_d(T)$ there is a region with ferrimagnetic long-range order, the model remains disordered for $J > J_d(T)$ down to $T = 0$.

Submitted to: *JSTAT*



Figure 1. Anisotropic kagome model (left) and auxiliary model (right). The factors of the transfer matrix acting in and between the different rows are indicated on the right of the auxiliary model.

1. Introduction

Recent experimental work on the mineral volborthite has stimulated several theoretical investigations of the Heisenberg antiferromgnet on the spatially anisotropic kagome lattice, figure 1, [1, 2, 3]. In this model, the exchange coupling in one of the three lattice directions (J) differs from the couplings in the two other directions (J').

The spin ensemble is thus divided into a set of parallel chains of spins with coupling J and a set of middle spins, which are coupled to the chain spins by the exchange constant J'. The results for the ground state of this model, obtained in the studies [1, 2, 3], differ substantially from each other, particularly in the quasi-one-dimensional limit $J \gg J'$ where the spin chains are only weakly coupled to the middle spins .

In this article, motivated by the difficulties encountered in approximate treatments of the Heisenberg model, we address the Ising model with anisotropic couplings on the kagome lattice (AIK). As was first demonstrated by Kano and Naya [4], this model can be solved exactly. Its thermodynamic properties are discussed in detail in recent work by Wei Li et al. [5]. For $J \neq J'$, the ground states are immediately clear. In the region $J < J'$, the ground state is ferrimagnetically ordered, i. e. the spins on the chains are ordered ferromagnetically, and so are the middle spins, but their direction is antiparallel to the chain spins. For $J > J'$, the chain spins are ideally ordered antiferromagnetically in the ground state leaving the middle spins completely uncoupled. As a consequence, the chains are also decoupled from one another, so that the spins of each chain can form either one of the two antiferromagnetic orders. A simple counting argument yields a rest entropy of $\mathcal{S}^{rest}_{aniso} = \frac{1}{3} \ln 2 \simeq 0.231$ per spin in this case. For the isotropic model, $J = J'$, the value of the rest entropy does not follow from a simple counting argument. Kano and Naya [4] find in this case

$$
\begin{aligned}
\mathcal{S}_{iso}^{rest} &= \frac{1}{24\pi^2}\int_0^{2\pi}\int_0^{2\pi} \ln\left[21-4(\cos(\omega_1)+\cos(\omega_2)+\cos(\omega_1+\omega_2))\right]d\omega_1 d\omega_2 \\
&\simeq 0.5018\ . \qquad (1)
\end{aligned}
$$

In the present paper, we shall reformulate the transfer matrix formalism of Kano and Naya with the aim to obtain not only the partition function, but also the correlation function $\chi(m,n)$ between two middle spins in the same row (see figure 1). We expect this to be the most interesting of the various spin-spin correlation functions of the AIK model.

The article is organised as follows. In section 2, we consider a slightly modified model that has been devised by Kano and Naya to allow for the application of the transfer matrix method, and that reproduces the AIK model in a certain limit. We present the transfer matrix, discuss its symmetry properties and convert it into a product of Grassmann functionals [6]. In section 3, we discuss the structure of the eigenvectors of the transfer matrix and calculate its eigenvalues. The latter are then used to construct the phase diagram of the AIK model as far as this is possible without knowledge of the correlation function. Section 4 is the central part of this article; it is devoted to our study of the above mentioned two-spin correlation. As in numerous previous investigations of correlations in Ising models on various two-dimensional lattices [7, 8, 9, 10, 11, 12, 13, 14, 15], the correlation function considered here is found to be represented rigorously by a Toeplitz determinant. By evaluating this determinant [16, 17] we obtain analytic expressions for $\chi(m,n)$ in the limit of large distances between the spins, $|n-m| \gg 1$, in various regions and on various special lines of the phase diagram. We support our asymptotic results by exact numerical evaluations of the Toeplitz determinant, which yield the correlation function also for short distances. In section 5 we present and discuss our results. Technicalities of the calculations that lead to these results are contained in appendices A, B, C. In appendix D, we present a simple perturbative calculation, which is valid in the limit of large chain coupling, $J/J' \gg 1$, and which qualitatively reproduces the behaviour of the correlation function in this limit.

After completion of this work, we became aware of a publication by M. B. Geilikman [18], which addresses the same topic as the present paper. Using the methods of Vdovichenko [19, 20], Geilikman arrives at the same Toeplitz determinant as we, to represent the spin-spin correlation function. However, his evaluation of this determinant yields results that differ qualitatively from our expressions for $\chi(m,n)$ in the most interesting part of the phase diagram.

2. Transfer matrices

Following Kano and Naya [4], we split each intermediate spin of the AIK into two Ising variables σ_j, σ_{j+1}, which are coupled ferromagnetically with a coupling l. At some point of our calculation we take the limit $l \to \infty$ so that these two Ising spins freeze into the original intermediate spin. The auxiliary model, which results from the splitting, is shown in the right part of figure 1. Obviously, it is periodic with period 4 in the (vertical) transfer direction. We consider a lattice of M (M divisible by 4) rows with N (N even) sites in each row and impose periodic boundary conditions in both lattice directions. For the auxiliary model, the transfer matrix, whose highest eigenvalue yields the free energy of this model, is given by (see figure 1)

$$T_0\, T_2\, T_0\, T_1\, T_0\, T_3\, T_0\, T_1, \tag{2}$$

where

$$T_0 = (e^{2K'} - e^{-2K'})^{N/2} \prod_{n=0}^{N-1} \sigma_n^x \quad e^{a \sum_{n=0}^{N-1} \sigma_n^x}, \tag{3}$$

$$T_1 = e^{-K \sum_{n=0}^{N-1} \sigma_n^z \sigma_{n+1}^z}, \tag{4}$$

$$T_2 = e^{\ell \sum_{n=0}^{(N-2)/2} (\sigma_{2n+1}^z \sigma_{2n+2}^z - 1)}, \tag{5}$$

$$T_3 = e^{\ell \sum_{n=0}^{(N-2)/2} (\sigma_{2n}^z \sigma_{2n+1}^z - 1)}, \tag{6}$$

with

$$K = J/T \quad , \quad K' = J'/T \quad \text{and} \quad \tanh(a) = e^{-2K'}. \tag{7}$$

(Boltzmann's constant k_B is set equal to unity throughout this paper.)
Instead of K and K' we shall use

$$\Psi = \tanh(K) \quad \text{and} \quad \Phi = \tanh(K') \tag{8}$$

as variables in the sequel.

The expression (2) is only one of various possibilities to represent the transfer matrix of the auxiliary model. In fact, it is an inconvenient representation, since it is not symmetric. Instead we can work with the following matrix

$$T_2^{1/2}\, T_0\, T_1\, T_0\, T_3^{1/2} \;\cdot\; T_3^{1/2}\, T_0\, T_1\, T_0\, T_2^{1/2}, \tag{9}$$

which in turn is equivalent to

$$T = T_2\, T_0\, T_1\, T_0\, T_3 \;\cdot\; T_3\, T_0\, T_1\, T_0\, T_2 \tag{10}$$

in the limit $l \to \infty$, since T_2 and T_3 become projectors in this limit, $T_i^2 = T_i,\ i = 2, 3$. T is real and symmetric since $T_3 = s\, T_2\, s$, where s shifts all spin variables by one lattice unit, $s\, \sigma_n^x\, s = \sigma_{n+1}^x,\ \ s\, \sigma_n^z\, s = \sigma_{n+1}^z$, and therefore commutes with T_0 and T_1.

In the sequel, we follow the transfer-matrix method of Schulz, Mattis and Lieb [9]. The necessary calculations are straight-forward. However, since the kagome lattice and the auxiliary lattice are non-Bravais lattices, some pecularities arise in comparison to the square lattice case. Therefore, we sketch the transfer-matrix procedure briefly here.

In order to exhibit the free fermion nature of the AIK model, one transforms the expressions in the exponents of T_0, ... T_3 into quadratic forms in fermion operators b_n, $b_n^\dagger$ by applying the Jordan-Wigner transform [21]:

$$\sigma_n^x = 2b_n^\dagger b_n - 1, \tag{11}$$

$$\sigma_0^z = b_0^\dagger + b_0 \quad \text{and} \quad \sigma_n^z = (b_n^\dagger + b_n)\; e^{i\pi\sum_{j=0}^{n-1} b_j^\dagger b_j}, \;\; 0 < n < N. \tag{12}$$

This yields

$$T_0 = \left(\frac{4\Phi}{1-\Phi^2}\right)^{N/2} \; e^{-Na} \;\; e^{(2a+i\pi)\sum_{n=0}^{N-1} b_n^\dagger b_n}, \tag{13}$$

$$T_1 = e^{-K\sum_{n=0}^{N-1}(b_n^\dagger - b_n)(b_{n+1}^\dagger + b_{n+1})}, \tag{14}$$

$$T_2 = e^{\ell\sum_{n=0}^{(N-2)/2}\left[(b_{2n+1}^\dagger - b_{2n+1})(b_{2n+2}^\dagger + b_{2n+2}) - 1\right]}, \tag{15}$$

$$T_3 = e^{\ell\sum_{n=0}^{(N-2)/2}\left[(b_{2n}^\dagger - b_{2n})(b_{2n+1}^\dagger + b_{2n+1}) - 1\right]}, \tag{16}$$

or, after Fourier transformation to momentum space

$$\begin{aligned} T_0 \;&= \left[\frac{16\Phi^4}{(1-\Phi^2)^2}\right]^{N/4} \; e^{(2a+i\pi)\,\hat{h}_0}, \\ T_1 \;&= e^{-K\,\hat{h}_1}, \\ T_{2,3} &= e^{-2\ell\, N/4} \;\; e^{\frac{1}{2}\ell\,\hat{h}_{2,3}} \end{aligned} \tag{17}$$

with

$$\hat{h}_0 \;= \sum_{-\pi\le k<\pi} b_k^\dagger b_k \;, \tag{18}$$

$$\hat{h}_1 \;= \sum_{-\pi\le k<\pi} \Big[i\sin(k)\left(b_k^\dagger b_{-k}^\dagger + b_k b_{-k}\right) + \cos(k)\left(b_k^\dagger b_k - b_k b_k^\dagger\right)\Big] \tag{19}$$

$$\begin{aligned} \hat{h}_{2,3} = \sum_{-\pi\le k<\pi} &\Big[i\sin(k)\left(b_k^\dagger b_{-k}^\dagger + b_k b_{-k} \mp b_k^\dagger b_{k+\pi} \mp b_k b_{k+\pi}^\dagger\right) \\ &+ \cos(k)\left(b_k^\dagger b_k - b_k b_k^\dagger \pm b_k b_{\pi-k} \mp b_k^\dagger b_{\pi-k}^\dagger\right)\Big]. \end{aligned} \tag{20}$$

We work with periodic boundary conditions. For states with even (odd) numbers of fermions this requires that $b_N^\dagger = b_0^\dagger$ ($b_N^\dagger = -b_0^\dagger$) (cf. reference [9]). To satisfy these conditions one chooses in $b_n = \frac{1}{\sqrt{N}}\sum_k e^{ikn}\, b_k$ wave numbers $k = \frac{2\pi}{N}m$ or $k = \frac{2\pi}{N}(m+1/2)$, m integer, for the even and odd states, respectively.

The non-Bravais structure of the auxiliary lattice in figure 1 manifests itself in the structure of the hamiltonians $\hat{h}_{2,3}$: they couple not only fermionic states with wave numbers k and $-k$ as in the square lattice case, but also states with wave numbers k and $\pm k + \pi$. To deal with this situation we divide the Brillouin zone $-\pi \le k < \pi$ into 4 subintervalls, $-\pi \le k < -\pi/2$, $-\pi/2 \le k < 0$; $0 \le k < \pi/2$, $\pi/2 \le k < \pi$ and choose N to be a multiple of 4. Then, $\hat{h}_0$, $\hat{h}_1$ and $\hat{h}_{2,3}$ are diagonal in q and can be represented in the reduced Brillouin zone $0 \le q < \pi/2$. For instance,

$$\begin{aligned}
\hat{h}_0 &= \sum_{0<q<\pi/2} \hat{h}_{0q} + \hat{h}_{0b}^{even} \qquad \text{with} \\
\hat{h}_{0q} &= b^\dagger_{-\pi+q} b_{-\pi+q} + b^\dagger_{-q} b_{-q} + b^\dagger_q b_q + b^\dagger_{\pi-q} b_{\pi-q} \ , \ \text{and} \\
\hat{h}_{0b}^{even} &= b^\dagger_{-\pi} b_{-\pi} + b^\dagger_{-\pi/2} b_{-\pi/2} + b^\dagger_0 b_0 + b^\dagger_{\pi/2} b_{\pi/2} \ .
\end{aligned} \tag{21}$$

The extra term $\hat{h}_{0b}^{even}$ corresponds to the values $k = -\pi,\ -\pi/2,\ 0, \pi/2$ of the original Brillouin zone. It occurs only if $\hat{h}_0$ acts in the space of *even states*. Then,

$$T_0^{even} = \left(\frac{16\Phi^4}{(1-\Phi^2)^2} \right)^{N/4} T_{0b} \prod_{0<q<\pi/2} T_{0q}, \tag{22}$$

where $T_{0b} = e^{2a\hat{h}_{0b}^{even}}$ and $T_{0q} = e^{2a\hat{h}_{0q}}$. Similarly, T_1 and $T_{2,3}$ are products of matrices acting in the subspaces of the individual wave numbers q. We remark that while these products consist of $N/4 - 1$ factors corresponding to the wavenumbers $0 < q < \pi/2$, there is only 1 factor of the type T_{0b} in each of the matrices T_0, T_1, T_2, and T_3. In the present paper we are always interested in the thermodynamic limit $N \to \infty$ of our results. In this limit the factors T_{0b} play no role and will therefore be neglected in the sequel.

3. Eigenvalues and eigenvectors of the symmetric transfer matrix

Our next aim is to determine the largest eigenvalue and the corresponding eigenfunction of the symmetric transfer matrix (10). It consists of factors that are all exponentials of quadratic forms of fermion operators. Hence T is also an exponential of a quadratic form of fermion operators. To determine this last exponential, the multiplication of the individual factors in equation 10 has to be performed explicitly. To achieve this, we find it convenient to work with the representation of the fermion operators b_q, $b_q^\dagger$ by Grassmann variables γ_q, γ_q^*. In appendix A we briefly describe the steps that are necessary to convert operators such as $T_0, 1, 2, 3$ into the so called *matrix form*. For details, we refer the reader to the book by F. A. Berezin [22], in particular to chapter 6. We introduce the shorthand notation $\vec{\gamma}_q = \{\gamma_q, \gamma_{-q}, \gamma_{-\pi+q}, \gamma_{\pi-q}\} \equiv \{\gamma_1, \gamma_2, \gamma_3, \gamma_4\}$. (There is no need to distinguish the row vector $\vec{\gamma}_q$ from its adjoint column vector $\vec{\gamma}_q^{\,t}$ here, since dyadic products $\vec{\gamma}_q \cdot \vec{\gamma}_q^{\,t}$ will not occur.) In matrix form, T_0, T_1 and $T_{2,3}$ read (here and in the sequel we use the shorthand notation $\cos(q) = c_q$, $\sin(q) = s_q$)

$$\tilde{T}_0(\gamma^*, \gamma) = \left(\frac{4\Phi^2}{1-\Phi^2} \right)^2 e^{-h_0 \, \vec{\gamma}_q^{\,*} \vec{\gamma}_q} \ , \quad h_0 = \frac{1}{\Phi}, \tag{23}$$

$$\tilde{T}_1(\gamma^*, \gamma) = \frac{(1+\Psi^2)^2 - 4c_q^2 \Psi^2}{(1-\Psi^2)^2} \ e^{\vec{\gamma}_q^{\,*} \, h_1 \, \vec{\gamma}_q \, + \, \frac{1}{2} \vec{\gamma}_q^{\,*} \, h_1' \, \vec{\gamma}_q^{\,*} \, - \, \frac{1}{2} \vec{\gamma}_q \, h_1'^{*} \, \vec{\gamma}_q} \tag{24}$$

with

$$h_1 = \begin{pmatrix} \frac{1-\Psi^2}{1+\Psi^2+2c_q\Psi} & 0 \\ 0 & \frac{1-\Psi^2}{1+\Psi^2-2c_q\Psi} \end{pmatrix}, \quad h_1' = \frac{2s_q\Psi}{1-\Psi^2}\, h_1 \begin{pmatrix} \sigma^y & 0 \\ 0 & -\sigma^y \end{pmatrix}, \tag{25}$$

$$\tilde{T}_2(\gamma^*,\gamma) = \frac{1}{4}\, e^{\vec{\gamma}_q^*\,(1+h_2)\,\vec{\gamma}_q \;+\; \frac{1}{2}\vec{\gamma}_q^*\, h_2'\, \vec{\gamma}_q^* \;-\; \frac{1}{2}\vec{\gamma}_q\, h_2'^*\, \vec{\gamma}_q} \tag{26}$$

with

$$h_2 = \begin{pmatrix} c_q & is_q\sigma^z \\ -is_q\sigma^z & -c_q \end{pmatrix}, \quad h_2' = -\begin{pmatrix} s_q & ic_q\sigma^z \\ ic_q\sigma^z & s_q \end{pmatrix} \begin{pmatrix} \sigma^y & 0 \\ 0 & -\sigma^y \end{pmatrix}. \tag{27}$$

Here the limit $\ell \to \infty$ has already been taken.

From the remarks after equation (10) of section 2 it follows that

$$\begin{aligned} T &= T_{1/2} \cdot T_{1/2}, \quad \text{where} \\ T_{1/2} &= T_2\, T_0\, T_1\, T_0\, s\, T_2. \end{aligned} \tag{28}$$

Thus it suffices to determine the eigenvalues and the eigenstates of $T_{1/2}$. Following the technique described in appendix A, we determine the *matrix form* of $T_{1/2}$:

$$\begin{aligned} \widetilde{T_{1/2\,q}}(\gamma^*\gamma) &= \int d\lambda^* d\lambda\; d\mu^* d\mu\; d\nu^* d\nu\; d\rho^* d\rho\;\; e^{-\vec{\lambda}^*\vec{\lambda}}\; e^{-\vec{\mu}^*\vec{\mu}}\; e^{-\vec{\nu}^*\vec{\nu}}\; e^{-\vec{\rho}^*\vec{\rho}} \\ &\qquad \tilde{T}_2(\gamma^*,\lambda)\; \tilde{T}_0(\lambda^*,\mu)\; \tilde{T}_1(\mu^*,\nu)\; \tilde{T}_0(\nu^*,\rho)\; \tilde{T}_2(s\rho^*,\gamma) \\ &= v\; e^{\vec{\gamma}_q^*\, H\, \vec{\gamma}_q \;+\; \frac{1}{2}\vec{\gamma}_q^*\, H'\, \vec{\gamma}_q^* \;-\; \frac{1}{2}\vec{\gamma}_q\, H'^*\, \vec{\gamma}_q} \qquad (29) \\ \widetilde{T_{1/2}} &= \prod_q \widetilde{T_{1/2\,q}}\,. \qquad (30) \end{aligned}$$

(We have suppressed the momentum label q in the first two lines of this equation.) The shift operator acts on ρ^* as $s\,\rho^* = \{\rho_1^*,\ \rho_2^*,\ -\rho_3^*,\ -\rho_4^*\}$.

We point out here that being quadratic in fermion variables, the exponent in $\widetilde{T_{1/2\,q}}$, equation (29), is the hamiltonian of a system of noninteracting fermions.

The general structure of the hamiltonians H and H' derives from the structure of the hamiltonians h_0, h_1, h_1', h_2 and h_2', equations (23), (25), (27): h_0, h_1, h_2 and h_1', h_2' are of the form

$$\mathfrak{M} = \begin{pmatrix} a & ib\,\sigma^z \\ ic\,\sigma^z & d \end{pmatrix} \;\text{and}\;\; \mathfrak{M}\cdot u, \;\; u = \begin{pmatrix} \sigma^y & \\ & -\sigma^y \end{pmatrix}, \text{respectively}, \tag{31}$$

where a, b, c, and d are real scalars. Products of $\mathfrak{M}$ are again of the form $\mathfrak{M}$ and furthermore $[\mathfrak{M}, u] = 0$, $u^2 = 1$. Hence, in equation (29), H is of the form $\mathfrak{M}$ and H' is of the form $\mathfrak{M}\;u$. Furthermore, since $\widetilde{T_{1/2}}$ is a real symmetric quadratic form in the space of the Grassmann vectors $\vec{\gamma}_q$, H must be symmetric and H' must be antisymmetric. Hence H and H' must have the following structure:

$$H = \begin{pmatrix} a & ib\,\sigma^z \\ -ib\,\sigma^z & d \end{pmatrix} \quad \text{and} \quad H' = \begin{pmatrix} a' & ib'\,\sigma^z \\ ib'\,\sigma^z & d \end{pmatrix} \begin{pmatrix} \sigma^y & \\ & -\sigma^y \end{pmatrix}. \tag{32}$$

The real scalars a, b, d and a', b', d' are rational functions of Ψ and Φ. Explicit expressions for these quantities are given in appendix B. Their calculation is straight-forward.

Turning to the eigenvalue problem for the transfer matrix we first consider the space of even states. As an even eigenstate we choose in Grassmann representation

$$\begin{aligned}\phi_g(\gamma^*) = \phi_0 - i\phi_1\,\gamma_1^*\gamma_2^* + i\phi_2\,\gamma_3^*\gamma_4^* - \phi_3\,\frac{1}{\sqrt{2}}(\gamma_1^*\gamma_4^* + \gamma_2^*\gamma_3^*) \\ + \phi_4\,\gamma_1^*\gamma_2^*\gamma_3^*\gamma_4^*\,. \end{aligned} \tag{33}$$

Again the momentum variable q is suppressed here. Only products of Grassmann variables with total momentum $q = 0 \bmod \pi$ are included in this state. The coefficients $\phi_{0,1,2,3,4}$ of the eigenstate (33) are calculated from

$$\int d\lambda^* d\lambda\ \widetilde{T_{1/2}}(\gamma^*, \lambda)\ e^{-\lambda^*\lambda}\ \phi_g(\lambda^*) = \sqrt{E_q^{(g)}}\ \phi_g(\gamma^*), \tag{34}$$

where (see (32))

$$\begin{aligned}\widetilde{T_{1/2}}(\gamma^*, \lambda) = v\exp\{&a(\gamma_1^*\lambda_1 + \gamma_2^*\lambda_2) + d(\gamma_3^*\lambda_3 + \gamma_4^*\lambda_4) \\ &+ ib(\gamma_1^*\lambda_3 - \gamma_2^*\lambda_4 - \gamma_3^*\lambda_1 + \gamma_4^*\lambda_2) \\ &- ia'\gamma_1^*\gamma_2^* + id'\gamma_3^*\gamma_4^* - b'(\gamma_1^*\gamma_4^* + \gamma_2^*\gamma_3^*) \\ &- ia'\lambda_1\lambda_2 + id'\lambda_3\lambda_4 + b'(\lambda_1\lambda_4 + \lambda_2\lambda_3)\},\end{aligned} \tag{35}$$

and we observe that the products with nonzero (mod π) momenta, $\gamma_1^*\gamma_3^*$ and $\gamma_2^*\gamma_4^*$, are indeed not generated by the action of $\widetilde{T_{1/2}}$ on $\phi_g(\gamma^*)$.

Equation (34) results in

$$\mathcal{T}\begin{pmatrix}\phi_0\\ \phi_1\\ \phi_2\\ \phi_3\\ \phi_4\end{pmatrix} = \sqrt{E_q^{(g)}}\begin{pmatrix}\phi_0\\ \phi_1\\ \phi_2\\ \phi_3\\ \phi_4\end{pmatrix}. \tag{36}$$

The 5×5 matrix $\mathcal{T}$ is presented in appendix B. Using the explicit expressions for the elements a, b, d, a', b', d' (see appendix B) we find

$$\det(\mathcal{T} - z) = -z^5 - 2c_5\,z^4 - c_4\,z^3, \tag{37}$$

with

$$\begin{aligned} c_5 = & -8\, c_q^2 \left[-\left(\frac{1-\Psi}{1+\Psi}\right)^2 \frac{1+6\Phi^2+\Phi^4}{(1-\Phi^2)^2} + 2\left(\frac{1+\Phi^2}{1-\Phi^2}\right)^2 - \left(\frac{1+\Psi}{1-\Psi}\right)^2 \right] \\ & -8 \left[\frac{1-\Psi}{1+\Psi}\left(\frac{1+\Phi^2}{1-\Phi^2}\right)^2 + \frac{1+\Psi}{1-\Psi} \right]^2 , \end{aligned} \tag{38}$$

$$c_4 = 2^{14}\, c_q^2 \left(\frac{1-\Psi}{1+\Psi}\right)^2 \frac{(1+\Phi^2)^2\, \Phi^4}{(1-\Phi^2)^6} . \tag{39}$$

c_5 is negativ and hence the largest eigenvalue in (36) is

$$\sqrt{E_q^{(g)}} = -c_5 + \sqrt{c_5^2 - c_4} . \tag{40}$$

As we need to know the largest eigenvalue of the transfer matrix for all positive values of J/J', we have to also consider the space of odd states. The appropriate ansatz for an odd eigenstate is

$$\begin{aligned} \phi_u(\gamma^*) = {} & \bar{\phi}_1\, \gamma_1^* + \bar{\phi}_2\, i\gamma_3^* + \bar{\phi}_3\, i\gamma_1^*\gamma_3^*\gamma_4^* + \bar{\phi}_4\, \gamma_1^*\gamma_2^*\gamma_3^* \\ & + \bar{\phi}_{-1}\, \gamma_2^* + \bar{\phi}_{-2}\, i\gamma_4^* + \bar{\phi}_{-3}\, i\gamma_2^*\gamma_3^*\gamma_4^* + \bar{\phi}_{-4}\, \gamma_1^*\gamma_2^*\gamma_4^* . \end{aligned} \tag{41}$$

Here the products of Grassmann variables in the first four terms have total momentum q mod π, while the products in the last four terms have total momentum $-q$ mod π. As the momentum q is conserved (mod π) in our model, the eigenvalue equation resulting from

$$\int d\lambda^* d\lambda\ \widetilde{T_{1/2}}(\gamma^*, \lambda)\ e^{-\lambda^*\lambda}\ \phi_u(\lambda^*) = \sqrt{E_q^{(u)}}\ \phi_u(\gamma^*) \tag{42}$$

is block diagonal,

$$\mathcal{T}_\pm \begin{pmatrix} \bar{\phi}_{\pm 1} \\ \bar{\phi}_{\pm 2} \\ \bar{\phi}_{\pm 3} \\ \bar{\phi}_{\pm 4} \end{pmatrix} = \sqrt{E_q^{(u\pm)}} \begin{pmatrix} \bar{\phi}_{\pm 1} \\ \bar{\phi}_{\pm 2} \\ \bar{\phi}_{\pm 3} \\ \bar{\phi}_{\pm 4} \end{pmatrix} . \tag{43}$$

Explicit expressions for the matrices $\mathcal{T}_+$, $\mathcal{T}_-$ are to be found in appendix B. The two largest eigenvalues $\sqrt{E_q^{(u+)}}$, $\sqrt{E_q^{(u-)}}$ turn out to be identical,

$$\sqrt{E_q^{(u+)}} = \sqrt{E_q^{(u-)}} = -c_4 . \tag{44}$$

As a first application of the results obtained so far in this section, we identify the transition line between ordered and disordered states of the AIK. As is shown in section VB of reference [9] a transition from a high-temperature disordered phase to an ordered state at low temperatures occurs when the even and the odd eigenstates become degenerate as the temperature is decreased. Thus we have to compare the eigenvalues

$E_q^g = (-c_5 + \sqrt{c_5^2 - c_4}\,)^2$ and $E_q^u = c_4$ as functions of the temperature T in the different regions of J/J'.
1) $J/J' < 1$: Using the explicit expressions for c_4 and c_5, equations (39), (38), one finds after some algebra that $E_q^g > E_q^u$ for all $q \neq 0$. For $q = 0$ E_q^g and E_q^u become equal if

$$\frac{1-\psi}{1+\psi} = \frac{1-\phi^2}{2\phi} \quad \text{or} \quad J = \frac{T_c}{2} \ln \frac{\sinh^2(\frac{2J'}{T_c})}{2\cosh(\frac{2J'}{T_c})}. \tag{45}$$

This defines the transition line between the ferrimagnetic and paramagnetic phase of the AIK (see figure 4 below) which was already determined by Kano and Naya.

2) $J = J'$, isotropic kagome model: Here $E_q^{(g)}$ is larger than $E_q^{(u)}$ for all momenta q and for all temperatures including $T = 0$. This reflects the fact that the ground state of the Ising model on the kagome lattice is disordered.

3) $J/J' > 1$. Using the explicit expressions for c_4 and c_5 again, one finds that $E_q^{(u)}/E_q^{(g)}$ attains its maximum at $q = 0$. As T decreases to zero, $E_q^{(u)}/E_q^{(g)} \to 0$ for $q \neq 0$. However, for $q = 0$, the ratio $E_q^{(u)}/E_q^{(g)} \to 1$ as $T \to 0$, implying that there is long range order in the ground state of the AIK.
These facts will be discussed in more detail below after we have determined the correlation function between the middle spins.
Since $E_q^{(g)} > E_q^{(u)}$ for all $q > 0$, the free energy per site of the AIK is

$$f = -\frac{4}{N}\, T \sum_{0 \le q < \pi/2} \ln(E_q^{(g)})\ . \tag{46}$$

In figure 2 we show the entropy per site that results from this expression.

Figure 2. Entropy $\mathcal{S}$ per site of the AIK model. J' has been set to unity here. Temperatures (from bottom to top): 0.05, 0.3, 0.5, 0.75, 1.0, 1.5, 2.5, 10.0 in units of J'. As $T \to \infty$, the entropy $\mathcal{S}$ reaches its limit $\ln 2$. As $T \to 0$, $\mathcal{S}$ decreases to 0 for $J < J'$ while a rest entropy of $\frac{1}{3}\ln 2$ remains for $J > J'$ (see introduction).



Obviously, the isotropic point $J = 1$ plays an exceptional role in the thermodynamics of the AIK.

4. Correlation function between middle spins

For a system with M rows of spins the correlation function between two middle spins at sites m and n $(m < n)$ of the same row is given by

$$\chi(m,n) = \frac{\mathrm{tr}\left\{ T_{1/2}^{2M} \; \sigma_m^z \sigma_n^z \right\}}{\mathrm{tr}\left\{ T_{1/2}^{2M} \right\}} \tag{47}$$

where $T_{1/2}$ is the transfer matrix (28) defined in section 3. If the state $|\mathfrak{Z}\rangle$ with maximum eigenvalue of the transfer matrix $T_{1/2}$ is non-degenerate this reduces to

$$\chi(m,n) = \frac{\langle \mathfrak{Z}| \; \sigma_m^z \sigma_n^z \, |\mathfrak{Z}\rangle}{\langle \mathfrak{Z}|\mathfrak{Z}\rangle} \tag{48}$$

in the thermodynamic limit $M \to \infty$. Again, the further development in this section follows closely the method devised by Schultz, Mattis and Lieb [9] in section IV of their work. Applying the Jordan-Wigner transformation, equation (12) to the product $\sigma_m^z \; \sigma_n^z$ we have

$$\sigma_m^z \; \sigma_n^z = i b_m^y b_n^x e^{i\pi \sum_{j=m+1}^{n-1} b_j^\dagger \, b_j}, \tag{49}$$

where

$$b_j^x = b_j^\dagger + b_j \quad \text{and} \quad i b_j^y = b_j^\dagger - b_j \; . \tag{50}$$

Expanding the exponential on the r. h. s. of equation (49) we obtain for the correlation function (48)

$$\chi(m,n) = \langle \phi_g | (i b_m^y) \, (b_{m+1}^x) \, (i b_{m+1}^y) \cdots (b_{n-1}^x) \, (i b_{n-1}^y) \, (b_n^x) | \phi_g \rangle / \langle \phi_g | \phi_g \rangle . \tag{51}$$

Here, $|\phi_g\rangle$ is the ground state of $-\ln T_{1/2}$ in the even subspace in the fermionic representation (c. f. equation (33),

$$\begin{aligned}
|\phi_g\rangle \quad &= \prod_{q=0}^{\pi/2} |\phi_g\rangle_q, \\
&\text{where} \\
|\phi_g\rangle_q \quad &= \phi_0(q) - i\phi_1(q) \; b_{1q}^\dagger b_{2q}^\dagger + i\phi_2(q) \; b_{3q}^\dagger b_{4q}^\dagger \\
&\quad - \phi_3(q) \; \frac{1}{\sqrt{2}} (b_{1q}^\dagger b_{4q}^\dagger + b_{2q}^\dagger b_{3q}^\dagger) + \phi_4(q) \; b_{1q}^\dagger b_{2q}^\dagger b_{3q}^\dagger b_{4q}^\dagger |0\rangle
\end{aligned} \tag{52}$$

$(|0\rangle :$ fermion vacuum).

The fermion operators $b_{1q}^\dagger$, $b_{2q}^\dagger$, $b_{3q}^\dagger$ and $b_{4q}^\dagger$ act in the intervals $[0, \pi/2[$, $[-\pi/2, 0[$, $[-\pi, -\pi/2[$ and $[\pi/2 2\pi[$, respectively. Our choice of the even ground state in equation

(51) implies that by evaluating this expression we will obtain the correlation function in the disordered region of the phase diagram where $|\phi_g\rangle$ is the ground state. Indeed, we expect to find the most interesting results in this region.

Since $|\phi_g\rangle$ is the ground state of a system of noninteracting fermions, cf. section 3, the expectation value in equation (51) can be evaluated by the use of Wick's theorem, which requires the knowledge of the contractions of all pairs of operators of the product in equation (51. In appendix B the coefficients $\phi_\nu(q)$, $\nu = 0, \cdots, 4$ are found to be real for all q. This suffices to show that

$$\langle\phi_g|b_j^x\ b_l^x|\phi_g\rangle = 0 \quad \text{and} \quad \langle\phi_g|b_j^y\ b_l^y|\phi_g\rangle = 0 \tag{53}$$

for all pairs of sites $j \neq l$. Denoting the remaining contractions by $G_{j,l+1}$,

$$G_{j,l+1} = \langle\phi_g|i\ b_j^y\ b_{l+1}^x|\phi_g\rangle / \langle\phi_g|\phi_g\rangle, \tag{54}$$

we can thus cast the result for the correlation function (51) into the form

$$\chi(m,n) = \begin{vmatrix} G_{m,m+1} & G_{m,m+2} & \cdots & G_{m,n-1} & G_{m,n} \\ G_{m+1,m+1} & G_{m+1,m+2} & \cdots & G_{m+1,n-1} & G_{m+1,n} \\ \cdot & & & & \cdot \\ \cdot & & & & \cdot \\ \cdot & & & & \cdot \\ \cdot & & & & \cdot \\ \cdot & & & & \cdot \\ \cdot & & & & \cdot \\ G_{n-1,m+1} & & \cdots & G_{n-2,n} & G_{n-1,n} \end{vmatrix} . \tag{55}$$

In appendix C we find that $G_{j,l} = 0$, if j and l are both even, or both odd, and that $G_{j,l} = \delta_{j+1,l}$, if j is odd. Furthermore, the remaining elements $G_{j,l}$ are found to depend only on the difference $l - j$. Hence, denoting

$$G_{2j,2l+1} = a_{l-j} \tag{56}$$

we get for the correlation function between middle spins at the sites 1 and $2n+1$

$$\chi(1,2n+1) = \begin{vmatrix} 1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & a_0 & 0 & a_1 & \cdots & 0 & a_{n-1} \\ 0 & 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & a_{-1} & 0 & a_0 & \cdots & 0 & a_{n-2} \\ 0 & 0 & 0 & 1 & \cdots & 0 & 0 \\ \cdot & & & & & & \cdot \\ \cdot & & & & & & \cdot \\ 0 & 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & a_{-n+1} & 0 & a_{-n+2} & \cdots & 0 & a_0 \end{vmatrix}$$

$$= \begin{vmatrix} a_0 & a_1 & \cdots & a_{n-1} \\ a_{-1} & a_0 & \cdots & a_{n-2} \\ \cdot & & \cdots & \\ \cdot & & \cdots & \\ \cdot & & \cdots & \\ a_{-n+1} & a_{-n+2} & \cdots & a_0 \end{vmatrix} . \tag{57}$$

Thus, for the AIK, the correlation function between the middle spins is represented by a Toeplitz determinant, just like the spin-spin correlation function of the square lattice Ising model, cf. reference [9].

In appendix C we show that

$$a_j = \oint_{|z|=1} \frac{dz}{2\pi i} \, \frac{1}{z} \, \frac{1}{z^j} \, \cdot \, \frac{p_2 + p_1 z^{-1} + p_0 z^{-2}}{|p_2 + p_1 z^{-1} + p_0 z^{-2}|}, \tag{58}$$

where

$$\begin{aligned} p_0 &= \left(1 - \Phi^2\right)^4 \, \Psi^2, \\ p_1 &= -\left(1 - \Phi^2\right)^2 \, \left[(1 - \Phi^2)^2 \, (1 + \Psi^4) + 4\Phi^2 \, (1 - \Psi) \, (1 - \Psi^3)\right], \\ p_2 &= \left[(1 - \Phi^2)^2 \, \Psi - 2\Phi^2 \, (1 - \Psi)^2\right]^2 . \end{aligned} \tag{59}$$

This expression for the elements of the Toeplitz determinant (57) agrees with the corresponding expression in reference [18] apart from a sign difference in p_1, which is probably due to a misprint in [18].

As is described in detail in Chapter XI of reference [16], for large n the Toeplitz determinant (57) can be evaluated analytically by applying a technique that was developed to solve Wiener-Hopf sum equations. We apply this technique here, but refrain from presenting any details apart from the remark that, as in the case of the square lattice Ising model, we have to deal with the case of *index* $\nu = -1$ here (see reference [16], p. 250). The result takes the form

$$\chi(1, 2n+1) = (-)^n \, R(n+1) \, x_n^{(n)} \tag{60}$$

Asymptotically,

$$(-)^n \, R(n) \sim - \left[\frac{(1 - z_B^2)(1 - z_A^{-2})}{(1 - z_B z_A^{-1})^2} \right]^{1/4} \tag{61}$$

and

$$x_n^{(n)} \sim - \oint_{|z|=1} \frac{dz}{2\pi i} \, z^{n-1} \, \sqrt{\frac{1 - z \, z_B}{1 - z \, z_A^{-1}}} \sqrt{\frac{1 - z^{-1} \, z_B}{1 - z^{-1} \, z_A^{-1}}} . \tag{62}$$

(Here and in the sequel the relation symbol " $\sim$ " implies that corrections to the right hand side of the expressions vanish in the limit $n \to \infty$.) The real quantities z_A and z_B are the roots of the the quadratic form $p_2 z^2 + p_1 z + p_0$ (see equation (58)),

$$z_{A,B} = -\frac{p_1}{2p_2} \pm \sqrt{\left(\frac{p_1}{2p_2}\right)^2 - \frac{p_0}{p_2}} \,. \tag{63}$$

Since $p_1 < 0$, the roots z_A and z_B are positive and $z_B \leq 1 \leq z_A$ for all values of the parameters Φ and Ψ outside the ordered region. However, depending on the values of $K = J/T$ and $K' = J'/T$ either ① $z_B < z_A^{-1}$ or ② $z_B > z_A^{-1}$.

① $z_B < z_A^{-1}$

We evaluate the integral in equation (62) by deforming the integration contour such that it encircles the branch cut of the integrand which extends from z_B to z_A^{-1},

$$-\,x_n^{(n)} \sim \int_{z_B}^{z_A^{-1}} \frac{dt}{\pi}\, t^{n-1} \sqrt{\frac{1 - t\, z_B}{1 - t\, z_A^{-1}}\, \frac{t - z_B}{z_A^{-1} - t}} \,. \tag{64}$$

In the limit of large n we obtain (see equation (60))

$$\chi(1, 2n+1) \sim \left[\frac{(1 - z_B^2)(1 - z_A z_B)^2}{1 - z_A^{-2}}\right]^{1/4} \cdot \sqrt{\frac{1}{\pi n}}\, z_A^{-n} \,. \tag{65}$$

② $z_A^{-1} < z_B$

Proceeding in the same manner as in case ① we find

$$-\,x_n^{(n)} \sim \int_{z_A^{-1}}^{z_B} \frac{dt}{\pi}\, t^{n-1} \sqrt{\frac{1 - t\, z_B}{1 - t\, z_A^{-1}}\, \frac{z_B - t}{t - z_A^{-1}}} \tag{66}$$

and hence in the limit of large n

$$\chi(1, 2n+1) \sim -\left[\frac{(1 - z_B^2)^3 (1 - z_A^{-2})}{(1 - z_B z_A^{-1})^4} \left(\frac{z_A z_B}{z_A z_B - 1}\right)^2\right]^{1/4} \cdot \sqrt{\frac{1}{4\pi n^3}}\, z_B^n \,. \tag{67}$$

We remark here that for the expressions (65) and (67) to be valid, the distances n must be large enough so that the condition

$$n \gg \bar{n} \equiv |\ln(z_A z_B)|^{-1} \tag{68}$$

is satisfied.

The plot of $\bar{n}$ in figure 3 indicates that in large regions of the phase diagram, in particular for $J > 1$, the correlation function $\chi(1, 2n+1)$ attains its asymptotic behaviour only for rather large n.



Figure 3. $\bar{n}$ ($J' = 1$); as $J \to 0.91$, $\bar{n} \to 0$ since $z_A z_B \to 0$ in this limit, cf. equation (68).

5. Results and discussion

The most remarkable feature of the phase diagram, figure 4, is the disorder line $J_d(T)$, which separates the regions ① and ② defined in the previous section from each other, see reference [23] and references therein. It is determined by the equation

$$\Phi^2 = (\tanh(J'/T))^2 = \tanh(J/T) = \Psi \ . \tag{69}$$

Figure 4. Phase diagram (T in units of J', J' is set to unity). FM: ferrimagnetically ordered region, full line: critical line $J_c(T)$; ①, ② disordered regions, separated by the disorder line $J_d(T)$ (dashed line).



On this line

$$\frac{p_2 + p_1 z^{-1} + p_0 z^{-2}}{|p_2 + p_1 z^{-1} + p_0 z^{-2}|} = \frac{1}{z}. \qquad (|z| = 1) \tag{70}$$

Then it follows from Eq. (58) that in this case $a_j = \delta_{j,-1}$ and hence $\chi(1, 2n+1) = 0$, *i.e.* the middle spins are uncorrelated on the disorder line.

5.1. Unfrustrated Ising like region

In region ①, the asymptotic behaviour of the correlation function is given by

$$\chi(1, 2n+1) \sim C_1 \cdot \frac{e^{-\frac{2n}{\xi_1}}}{\sqrt{n}}, \quad \xi_1^{-1} = \frac{\ln(z_A)}{2}, \ \ C_1 > 0 \text{ (cf. equation (65)).} \tag{71}$$

This suggests that in this region the AIK model behaves like a unfrustrated Ising magnet in the disordered phase, cf. reference [16], Chapter XI. In fact, for $J = 0$ the AIK model, cf. figure 1, reduces to the antiferromagnetic Ising model on the square lattice whose edges are decorated by the chain spins, which are decoupled from each other. Tracing out these spins yields an effective ferromagnetic coupling between the middle spins with coupling constant $K_\Box = -\frac{1}{2} \ln\cosh(2K')$. Furthermore, for $J = 0$ equations (63) and (59) yield

$$z_B = 0 \quad \text{and} \quad z_A^{-1} = -\frac{p_2}{p_1} = \sinh^2(2K_\Box). \tag{72}$$

Then, the expression (65) for the correlation function between the middle spin of the AIK model reduces to the known expression for the correlation function between spins along one of the diagonals of the ensuing square model in the disordered region, cf. reference [16], Chapter XI.2.

To check the quality of the asymptotic expression (65) for the correlation function $\chi(1, 2n+1)$ we have calculated the determinant in equation (57) numerically. In figure 5 the asymptotic result (65) and the numerical result are compared after the exponential factor e^{-2n/ξ_1} has been divided out from both results. As is expected on account of the condition (68), for small and moderate distances $2n$ the agreement is worse if J is close to the disorder line where $\bar{n} \to \infty$.

On the critical line $T_c(J)$, cf. equation (45), we find $z_A = 1$ and

$$a_j = -\oint\limits_{|z|=1} \frac{dz}{2\pi i}\, \frac{1}{z}\, \frac{1}{z^j} \cdot \sqrt{\frac{-1}{z^2}\, \frac{z - z_B}{1 - z\, z_B}}\,. \tag{73}$$

This is very similar to the corresponding expression for the square lattice Ising model, equation (5.1) in Chapter XI of reference [16]. Applying the same method as McCoy and Wu, we arrive at

$$\chi(2n+1, 1) \sim \left[(1 - z_B)^2 (1 - z_B^2)\right]^{(1/4)} \frac{A}{n^{(1/4)}} \ \ (A : \text{see reference [16]}) \tag{74}$$



Figure 5. Comparison of the asymptotic (line) and the numerical results (symbols) for $\chi(1,2n+1)$ for $J=0.7, \bar{n}=0.3$ (left) and $J=0.825, \bar{n}=17.5$ (right) after division by e^{-2n/ξ_1}, see text ($J'=1$).

where

$$z_B = \frac{(1+2\Phi^2-\Phi^4)^2(1-2\Phi^2-\Phi^4)^2}{(1+8\Phi^2+10\Phi^4+8\Phi^6+\Phi^8)^2}, \quad A = 0.645\cdots \quad . \tag{75}$$

In summary, the results presented here indicate clearly that in region ① the features of the AIK model differ only quantitatively from those of an unfrustrated Ising model with antiferromagnetic nearest neighbour interaction. The middle spins of one row of the model are correlated ferromagnetically. The appearance of a ferrimagnetically ordered region for small coupling J and low temperatures fully agrees with this picture of an unfrustrated model.

5.2. Chain dominated region

The features of the AIK model change qualitatively when the coupling J is increased across the disorderline into the region ②. The disordered phase extends down to $T=0$ in this region, and the correlation function of the middle spins changes from the expression in equation (71) to

$$\chi(1,2n+1) \sim -C_2 \cdot \frac{e^{-\frac{2n}{\xi_2}}}{\sqrt{n^3}}, \tag{76}$$

$$\text{with} \qquad \xi_2^{-1} = -\frac{\ln(z_B)}{2}, \quad C_2 > 0 \quad \text{(see equation (67))}.$$

This implies that all the middle spins in one row of the AIK model are predominantly correlated antiparallel to one another. In Appendix D we present a simple perturbative evaluation of the correlation function which is valid for $J' \ll J$ and $J'/T \ll 1$. In this limit, the correlations between the middle spins of one row are mediated by the spins of two adjacent chains which are correlated antiferromagnetically and to which the middle spins are weakly coupled. This mechanism leads quite naturally



Figure 6. Comparison of the asymptotic (line) and the numerical results (symbols) for $\chi(1, 2n+1)$ for $J = 0.91, \bar{n} = 0.1$ (left) and $J = 1.5, \bar{n} = 54.4$ (right) after division by e^{-2n/ξ_2} $(J' = 1)$.

to a negative correlation between the any pair of middle spins of the same row.

As we have mentioned in the Introduction, for $J > J'$ the middle spins decouple from the chain spins in the limit $T \to 0$ and become therefore completely uncorrelated. To see how this happens as the temperature is decreased to zero we expand z_A and z_B (cf. equation (63)) for low temperatures, i. e. $\epsilon = 2e^{-2J'/T} \ll 1, \quad \xi_2^{-1} = e^{-2(J-J')/T} \ll 1$,

$$\begin{aligned} z_A^{-1} &= 1 - 2\xi_2^{-1} + \frac{3}{2}\xi_2^{-2} + \mathcal{O}(\xi_2^{-1}\epsilon, \xi_2^{-2}\epsilon, \xi_2^{-3}), \\ z_B &= 1 - 2\xi_2^{-1} + \frac{5}{2}\xi_2^{-2} + \mathcal{O}(\xi_2^{-1}\epsilon, \xi_2^{-2}\epsilon, \xi_2^{-3}). \end{aligned} \tag{77}$$

With these expansions, equation (67) yields

$$\chi(1, 2n+1) \sim -\frac{1}{2}\, \xi_2^{-2}\, e^{-2n/\xi_2}\ . \tag{78}$$

Although the correlation length ξ_2 tends to infinity as $T \to 0$, $\chi(1, 2n+1)$ vanishes in this limit since the prefactor ξ_2^{-2} vanishes.

To compare the asymptotic result (67) for $\chi(1, 2n+1)$ in the parameter region ② with the exact result (57) we have evaluated the latter numerically as in the other regime. The comparison is shown in figure 6. As in region ①, the agreement between the asymptotic and the exact results becomes worse for small and moderate distances n as $\bar{n}$ increases.

5.3. The point $J' = J = 1$, $T = 0$

From our discussion of the rest entropy of the AIK model in the introduction, it follows that in the limit $T \to 0$ this model behaves irregularly at the isotropic point $J' = J = 1$. Here, we wish to discuss the behaviour of the correlation function between the middle spins as the isotropic point is approached from different directions in the phase diagram. We fix a direction by choosing a fixed value for the parameter α in the equation

$$J - J' = \alpha T, \tag{79}$$

which represents straight lines with different slopes in the phase diagram that intersect at the isotropic point. For instance, $\alpha = -\ln 2$ is the slope of the phase boundary $T = T_c(J - J')$ at $J' = J = 1$, $\alpha = -\frac{1}{2}\ln 2$ is the slope of the disorder line at this point and $\alpha = 0$ is the line perpendicular to the abscissa in $J' = J = 1$. By parametrising

$$\Phi = 1 - \epsilon \qquad \Psi = 1 - \epsilon\sqrt{s} \quad \text{where} \quad s = e^{-4\alpha} \tag{80}$$

we obtain from equations (59), (63)

$$z_A^{-1} = z_B \left[\frac{2-s}{2}\right]^2, \quad z_B = \frac{4+6s}{(2-s)^2} - \sqrt{\left[\frac{4+6s}{(2-s)^2}\right]^2 - \left[\frac{2}{2-s}\right]^2} \quad . \tag{81}$$

This yields for the different cases:

$s = 16$:	on the phase boundary	χ as in equation (74),	$z_B = 1/49$
$16 > s > 4$:	in region ①	χ as in equations (65),(71)	$z_A^{-1} > z_B$
$s = 4$:	on the disorder line	$\chi = 0$	$z_A^{-1} = z_B = 1$
$4 > s > 0$:	in region ②	χ as in equations (67), (76)	$z_B > z_A^{-1}$
$s = 1$:	on the perpendicular	χ as in equation (76)	$\xi_2 \simeq 1.25$
$s = 0$:	on the line $T = 0$	$\chi = 0$ cf. equations (60), (61) and (62)	$z_A^{-1} = z_B = 1$ as on the disorder line

6. Summary and outlook

In this paper we have adapted the transfer matrix technique devised in reference [9] for the solution of the square lattice Ising model to the case of the Ising model on the anisotropic kagome model (AIK). The isotropic kagome model is fully frustrated, but as the coupling J in one of the three lattice directions is varied from zero to infinity, the model changes from the unfrustrated square lattice model to a set of antiferromagnetic chains. We have calculated the pair correlation function of middle spins, i. e. of spins on intermediate rows between the chains with coupling J, (i) asymptotically exact, (ii) numerically and for the case $J \gg 1$ (iii) by a perturbative method. We find that the phase diagram of the AIK model is divided into two sections by a disorder line $J_d(T)$. On this line, the correlation between the middle spins vanishes exactly. The following statements need to be corroborated by a calculation of the correlations between the chain spins, but this was not a subject of the present paper. For $J < J_c(T)$ the model is ferrimagnetically ordered: we expect the chain spins and the middle spins to be ordered ferromagnetically but antiparallel relative to each other; in the region $J_c(T) < J < J_d(T)$ we expect short range order of the same kind. (Short range ferromagnetic order between the middle spins is found in the present paper.) In summary, for $J < J_d(T)$ we expect the AIK model to behave like an unfrustrated Ising model.

At the disorder line, the correlation between the middle spins changes qualitatively. For $J_d(T) < J$ there is no long range order at any temperature. The three methods (i -

iii) show that the pair correlation function of the middle spins is negative throughout this region. It decays exponentially with a correlation length $\xi_2(T)$ that diverges in the limit $T \to 0$. However, as $T \to 0$ the amplitude of the correlation function becomes proportional to $\xi_2(T)^{-2}$, i. e. the middle spins become uncorrelated as $T \to 0$. Since the chains are antiferromagnetically ordered at $T = 0$, the loss of correlation between the middle spins was to be expected.

To complete the picture of the spin correlations of the AIK model various correlation functions in addition to the middle spin correlation function need to be calculated. We expect to be able to perform the necessary calculations by applying the transfer matrix technique devised here.

Acknowledgement

We thank R. Wynands for helpful comments on the manuscript of this article.

Appendix A. Working with Grassmann variables

Here we collect rules to be followed in transforming operators from their fermion representation, e. g. expressions (3) …(6) for the transfer matrices T_0 …T_3, into their representation by Grassmann variables and of the rules that were important in our manipulations of such operators. We adhere to the terminology of reference [22].

If $\hat{T}(b^\dagger, b)$ is a normal ordered fermionic operator ($b^\dagger$, b: fermion creation and annihilation operators), then $T(\gamma^*, \gamma) := \hat{T}\Big|_{b^\dagger\to\gamma^*,\, b\to\gamma}$, where γ are Grassmann numbers and γ^* their complex conjugates, is called the "normal form" of T, and $\tilde{T}(\gamma^*, \gamma) := T(\gamma^*, \gamma)\, e^{\gamma^*\gamma}$ is called the "matrix form" of T. ($b^\dagger$, b and γ^*, γ can be vectors of fermion operators or Grassmann numbers, respectively.)

Gaussian integral: For an arbitrary $N \times N$ matrix H we have

$$\prod_n \int d\gamma_n^* d\gamma_n \;\; e^{-\gamma_n^* H_{nm} \gamma_m} = \det H. \tag{A.1}$$

(In the exponent, automatic summation over the indices is implied.)

Gaussian integral with external sources: for an antisymmetric $2N \times 2N$ matrix R:

$$\int d\lambda^* d\lambda \; e^{\frac{1}{2}\begin{pmatrix} \lambda & \lambda^* \end{pmatrix} R^{-1} \begin{pmatrix} \lambda \\ \lambda^* \end{pmatrix} + \begin{pmatrix} \gamma & \gamma^* \end{pmatrix} \begin{pmatrix} \lambda \\ \lambda^* \end{pmatrix}}$$

$$= \pm\sqrt{\det R^{-1}} \;\; e^{\frac{1}{2}\begin{pmatrix} \gamma & \gamma^* \end{pmatrix} R \begin{pmatrix} \gamma \\ \gamma^* \end{pmatrix}} \tag{A.2}$$

Here, γ and λ are N-dimensional Grassmann vectors. The determinant of an antisymmetric matrix is always positiv. The sign of the square root depends on the order of the Grassmann integrations. It is irrelevant for our purposes.

Trace:

$$\mathrm{tr}\left\{\hat{T}\right\} = \int d\gamma d\gamma^* \, e^{\gamma^*\gamma} \, \tilde{T}(\gamma^*, \gamma) \tag{A.3}$$

.

Product of fermionic operators:

$$\hat{A} \cdot \hat{B} = \hat{C} : \quad \tilde{C}(\gamma^*\gamma) = \int d\lambda^* d\lambda \, \tilde{A}(\gamma^*\lambda)\tilde{B}(\lambda^*\gamma) \, e^{-\lambda^*\lambda}. \tag{A.4}$$

In our case all products in question can be evaluated explicitly since the integrals that have to be performed are gaussian (cf. equation (29)).

Appendix B. The matrices $\tilde{\mathcal{T}}$, $\tilde{\mathcal{T}}_\pm$

As the matrices $\mathcal{T}$, $\mathcal{T}_+$ and $\mathcal{T}_-$ play a central role in our analysis of the AIK model, we list their elements explicitely here in terms of our basic variables Ψ and Φ (cf. equation (8)). $\mathcal{T}$ is a real symmetric 5×5 matrix and $\mathcal{T}_\pm$ are real symmetric 4×4 matrices. Defining $\bar{\mathcal{T}}$ and $\bar{\mathcal{T}}_\pm$ by

$$\mathcal{T} = v \cdot \bar{\mathcal{T}}, \quad \mathcal{T}_{pm} = v \cdot \bar{\mathcal{T}}_{pm} \tag{B.1}$$

we have

$$\begin{aligned}
&\bar{\mathcal{T}}_{11} = 1, \quad \bar{\mathcal{T}}_{12} = a', \quad \bar{\mathcal{T}}_{13} = d', \quad \bar{\mathcal{T}}_{14} = \sqrt{2}b', \quad \bar{\mathcal{T}}_{15} = a'd' + b'^2, \\
&\bar{\mathcal{T}}_{22} = a^2 + a'^2, \quad \bar{\mathcal{T}}_{23} = a'd' - b^2, \quad \bar{\mathcal{T}}_{24} = \sqrt{2}(a'b' - ab), \\
&\bar{\mathcal{T}}_{25} = a'\left(a'd' + b'^2\right) - a'b^2 + d'a^2 - 2abb', \\
&\bar{\mathcal{T}}_{33} = d^2 + d'^2, \quad \bar{\mathcal{T}}_{34} = \sqrt{2}(bd + b'd'), \\
&\bar{\mathcal{T}}_{35} = d'\left(a'd' + b'^2\right) + a'd^2 - d'b^2 + 2bdb', \\
&\bar{\mathcal{T}}_{44} = ad + b^2 + 2b'^2, \quad \bar{\mathcal{T}}_{45} = \sqrt{2}\,\Big(b'\left(a'd' + b'^2\right) \\
&\qquad\qquad\qquad\qquad\qquad\qquad +a'bd - d'ab + b'(ad + b^2)\Big), \\
&\bar{\mathcal{T}}_{55} = \ \left(a'd' + b'^2\right)^2 + a'^2d^2 + d'^2a^2 \\
&\qquad\quad +4b'^2ad - 2a'd'b^2 + 4a'b'bd - 4b'd'ab
\end{aligned} \tag{B.2}$$

and

$$\begin{aligned}
&\bar{\mathcal{T}}_{\pm 11} = a, \quad \bar{\mathcal{T}}_{\pm 12} = -b, \quad \bar{\mathcal{T}}_{\pm 13} = ad' \mp bb', \quad \bar{\mathcal{T}}_{\pm 14} = \mp ab' - ba', \\
&\bar{\mathcal{T}}_{\pm 22} = d, \quad \bar{\mathcal{T}}_{\pm 23} = \pm db' - bd', \quad \bar{\mathcal{T}}_{\pm 24} = a'd \pm bb',
\end{aligned}$$

$$\begin{aligned}
&\bar{\mathcal{T}}_{\pm 33} = ad^2 \mp 2bb'd'^2 + db'^2, \quad \bar{\mathcal{T}}_{\pm 34} = \mp ab'd' \pm da'b' - b\left(a'd' + b'^2\right), \\
&\bar{\mathcal{T}}_{\pm 44} = ab'^2 + da'^2 \pm 2ba'b'.
\end{aligned} \tag{B.3}$$

In terms of the parameters Φ and Ψ the quantities v, a, d, b and a', d', b' are given by:

$$v = \frac{16\ (1 - {c_q}^2\Psi^2)\left(1 + \Phi^4\right)^2 - 4\ {s_q}^2\Phi^2\Psi\ (1 + \Phi^4) + 4\ (\Psi^2 - {c_q}^2)\,\Phi^4}{\left(1 - \Phi^2\right)^4\left(1 - \Psi^2\right)^2}\ ;$$

$$\begin{aligned}
&a = 2c_q\ (c_q + 1)\Phi^2(1 - \Phi^4)(1 - \Psi)^2(1 - \Psi^2)/den\ , \\
&d = -a|_{q \to \pi + q}\ , \\
&b = a\ \frac{s_q}{c_q + 1}
\end{aligned}$$

with

$$den = -(1 + \Phi^4 - 4\Phi^2\Psi + \Psi^2 + \phi^4\Psi^2)^2 + 4c_q^2(1 - \Phi^2\Psi)^2(\Phi^2 - \Psi^2)^2\ .$$

One sees immediately that $ad - b^2 = 0$. This identity has been used in establishing the matrices in equations (B.2) and (B.3).

$$\begin{aligned}
a' = \frac{s_q}{den}\Big(&((1 + \Phi^4)(1 + \Psi^2) - 4\Phi^2\Psi)^2 \\
&- 4c_q(1 - \Phi^2\Psi)(\Psi - \Phi^2)(\Phi^2(1 - \Psi)^2 + c_q\Psi(1 - \Phi^4))\Big)\ ,
\end{aligned}$$

$$d' = -a'|_{q \to q + \pi}\ ,$$

$$\begin{aligned}
b' = \frac{c_q}{den}(1 - \Phi^2)^2\Big(&(1 + \Psi^2)^2(1 + \Phi^4) + 2\Phi^2((1 - \Psi^2)^2 - 2\Psi - 2\Psi^3) \\
&- 4c_q^2\Psi(\Psi - \Phi^2)(1 - \Psi\Phi^2)\Big).
\end{aligned} \tag{B.4}$$

With this information explicit expressions for the eigenvalues and the eigenvectors of the transfer matrices that are needed in the main part of the paper can be obtained. We refrain from presenting the rather lengthy explicit expressions for the coefficients $\phi_1 \ldots \phi_4$ of the eigenvector (33) but we note certain symmetry properties of these coefficients that will be used to simplify expressions to be derived in the next appendix. First we observe that

$$\phi_4 = \phi_0, \tag{B.5}$$

and furthermore we find the following symmetries of ϕ_1, ϕ_3 and ϕ_3 as functions of q:

$$\begin{array}{r|c|c|c|}
 & \phi_1 \to & \phi_2 \to & \phi_3 \to \\
\hline
q \to -q & -\phi_1 & -\phi_2 & \phi_3 \\
q \to \pi + q & -\phi_2 & -\phi_1 & -\phi_3 \\
\hline
\end{array} \tag{B.6}$$

Appendix C. Calculation of the elements of the Toeplitz determinant

In this appendix we present the main steps of the calculations that lead to the result (58) for the elements a_j of the Toeplitz determinant (57) in the main text. We need to determine the expectation value (cf. equation (54))

$$\begin{aligned} G_{m,n} &= \frac{1}{\mathcal{N}_g}\langle\phi_g|ib^y_m b^x_n|\phi_g\rangle\ , \quad \text{where} \quad \mathcal{N}_g = \langle\phi_g|\phi_g\rangle \\ &= \frac{1}{\mathcal{N}_g}\langle\phi_g| - b_m\, b_n - (b_m\, b_n)^\dagger + b^\dagger_m\, b_n + (b^\dagger_m\, b_n)^\dagger\ |\phi_g\rangle(1-\delta_{m,n}) \\ &\quad + \left(\frac{1}{\mathcal{N}_g}\langle\phi_g|2\, b^\dagger_m b_m|\phi_g\rangle - 1\right)\ \delta_{m,n}\ , \end{aligned} \tag{C.1}$$

cf. equation (50). Let us first consider the expectation value

$$f_{m,n} = \frac{1}{\mathcal{N}_g}\langle\phi_g|b_m\, b_n|\phi_g\rangle. \tag{C.2}$$

With the representation

$$b_m = \frac{1}{\sqrt{N}}\sum_{0\le q<\pi/2}\left[e^{iqm}b_{1q} + e^{-iqm}b_{2q} + (-)^m e^{iqm}b_{3q} + (-)^m e^{-iqm}b_{4q}\right] \tag{C.3}$$

of the annihilation operator at site m by the momentum space operators $b_{1q}\ \ldots b_{4q}$ introduced in the main text (see paragraph below equation (52)) we find after a straightforward evaluation of expectation values of the type ${}_q\langle\phi_g|b_{\mu q}\, b_{\nu q}|\phi_g\rangle_q$, $(\mu,\ \nu = 1,\ldots,4)$

$$\begin{aligned} f_{m,n} =& \frac{1}{N}\sum_{0\le q<\pi/2}\frac{1}{\mathcal{N}_{gq}}\Big[-2\sin(q[m-n])\ (1-(-)^{m+n})\ \phi_0(\phi_1+\phi_2) \\ &\quad - 4\cos(q[m-n])\ ((-)^m-(-)^n)\ \frac{\phi_3}{\sqrt{2}}\phi_0\Big]\ . \end{aligned} \tag{C.4}$$

Here

$$\mathcal{N}_{gq} = {}_q\langle\phi_g|\phi_g\rangle_q = 2\phi_0^2 + \phi_1^2 + \phi_2^2 + \phi_3^2. \tag{C.5}$$

(The momentum label q has been suppressed on the r.h.s.) In equations (C.4) and (C.5) use has been made of the identity (B.5). Note that $f_{m,n}$ is real and hence

$$\frac{1}{\mathcal{N}_g}\langle\phi_g|b^\dagger_n\, b^\dagger_m|\phi_g\rangle = \frac{1}{\mathcal{N}_g}\langle\phi_g|b_m\, b_n|\phi_g\rangle = f_{m,n}\ .$$

Similarly,

$$g_{m,n} = \frac{1}{\mathcal{N}_g}\langle\phi_g|b^\dagger_m\, b_n|\phi_g\rangle \tag{C.6}$$

$$\begin{aligned} &= \frac{1}{N}\sum_{0\le q<\pi/2}\frac{1}{\mathcal{N}_{gq}}\Big[2\cos(q[m-n])\ \Big\{\phi_0^2+\phi_1^2+\frac{\phi_3^2}{2} \\ &\quad +(-)^{m+n}\ (\phi_0^2+\phi_2^2+\frac{\phi_3^2}{2})\Big\} \\ &\quad - 2\sin(q[m-n])\ \frac{\phi_3}{\sqrt{2}}\ [(-)^m-(-)^n]\ (\phi_1-\phi_2)\Big]\ . \end{aligned} \tag{C.7}$$

Obviously $g_{m,n}$ is real too, and with equation (C.5) it follows immediately that $g_{m,m} = 1/2$, so that the last term in the expression (C.1) vanishes. On account of the symmetries (B.6), the sums over q in Eqs. (C.4), (C.7) can be extended to the full Brillouin zone $-\pi \le q < \pi$,

$$\frac{1}{N} \sum_{0 \le q < \pi/2} \cdots \longrightarrow \frac{1}{4N} \sum_{-\pi \le q < \pi} \cdots \,.$$

Then,

$$\begin{aligned}
G_{m,n} &= 2\,(-f_{m,n} \;+\; g_{m,n})(1-\delta_{m,n}) \\
&= \frac{1}{2}[1-(-)^{n-m}]\,\gamma_{n-m} + \frac{1}{2}[(-)^m - (-)^n]\,\gamma'_{n-m}
\end{aligned}$$

with

$$\gamma_l \quad = \frac{4}{N} \sum_{-\pi \le q < \pi} \frac{1}{\mathcal{N}_{gq}} \frac{1}{4}(\phi_1 + \phi_2)\,[(\phi_1 - \phi_2)\cos(ql) - \;2\phi_0\;\sin(ql)]$$

and

$$\gamma'_l \quad = \frac{4}{N} \sum_{-\pi \le q < \pi} \frac{1}{\mathcal{N}_{gq}} \frac{\phi_3}{\sqrt{2}}\,[\phi_0\;\cos(ql) + \frac{1}{2}(\phi_1 - \phi_2)\;\sin(ql)]. \tag{C.8}$$

For the elements of the determinant (55) this yields

$$\begin{aligned}
G_{2m,2m'} &= 0\,, \qquad G_{2m+1,2m'+1} = 0 \\
G_{2m-1,2m'} &= \gamma_{2(m'-m)+1} - \gamma'_{2(m'-m)+1} \equiv a'_{m'-m} \\
G_{2m,2m'+1} &= \gamma_{2(m'-m)+1} + \gamma'_{2(m'-m)+1} \equiv a_{m'-m}\;.
\end{aligned} \tag{C.9}$$

Further simplifications of these expressions are possible on account of the following properties of the coefficients ϕ_1, ϕ_2 and ϕ_3 that were found by analysing the explicit expressions for these coefficients with computer algebra: They have a common denominator $\bar{D}_0(q)$ which is invariant under the operations $q \to -q$ and $q \to \pi + q$,

$$\phi_\nu(q) = \frac{\bar{D}_\nu(q)}{\bar{D}_0(q)}, \qquad \nu \;=\; 1, 2, 3. \tag{C.10}$$

Only two of the four coefficients $\bar{D}_0$, $\bar{D}_1$, $\bar{D}_2$ and $\bar{D}_3$ are independent. The relations between them are

$$-\frac{1}{\sqrt{2}} c_q \bar{D}_3(q) - \frac{1}{2} s_q [\bar{D}_1(q) + \bar{D}_2(q)] = \bar{D}_0(q)$$

and

$$-\frac{1}{\sqrt{2}} s_q \bar{D}_3(q) + \frac{1}{2} c_q [\bar{D}_1(q) + \bar{D}_2(q)] \;= \frac{1}{2}(\bar{D}_1(q) - \bar{D}_2(q)). \tag{C.11}$$

Using these relations one finds

$$a'_j = \delta_{j,0} \tag{C.12}$$

and, introducing $p = 2q$ as the momentum variable and denoting $\bar{D}_\nu(\frac{p}{2}) \equiv D_\nu$,

$$a_j = \frac{1}{N} \sum_{-\pi \le p < \pi} \left\{ \left[D_0^2 - \frac{1}{4}(D_1 - D_2)^2 \right] \cos(p[j+1]) \right.$$
$$\left. + D_0 \, (D_1 - D_2) \, \sin(p[j+1]) \right\} \frac{1}{D_0^2 + \frac{1}{4}(D_1 - D_2)^2} . \qquad \text{(C.13)}$$

In the continuum limit, $N \to \infty$, and with $z = e^{ip}$ as the integration variable, this leads to the expression (58) for a_j.

Appendix D. Perturbation expansion in J'/T

As we have mentioned in the Introduction, for $J > J'$ the chain spins of the AIK model are ideally ordered antiferrromagnetically in the ground state, and as a consequence the middle spins are completely decoupled from the chain spins. For weak interchain coupling $J' \ll J$ the order of the spins along the chains will prevail over a long distance for low but finite temperatures $T \ll J$, and the coupling between the chains will remain small as long as $J' \ll T$. In the sequel we will calculate the correlation function between the middle spins of row l of the kagome lattice, $\chi(2m+1, 2n+1) = \langle \, \sigma_{2m+1,l} \; \sigma_{2n+1,l} \, \rangle$, perturbatively with respect to J'. (In $\sigma_{j,l}$ the second subscript l denotes the row l of the lattice.) The hamiltonian of the AIK model couples the middle spins of row l to chain spins of the rows $l-1$ and $l+1$,

$$J' \, \sigma_{2m+1,l} \cdot [\sigma_{2m+1,l-1} + \sigma_{2m,l-1} + \sigma_{2m+1,l+1} + \sigma_{2m,l+1}] \; . \qquad \text{(D.1)}$$

In second order in J'/T one obtains for χ:

$$\begin{aligned}
&\chi(2m+1, 2n+1) \\
&= 2 \, (\frac{J'}{T})^2 \langle [\sigma_{2m+1,l+1} + \sigma_{2m,1}] \, [\sigma_{2n+1,l+1} + \sigma_{2n,1}] \rangle_{J'=0} + \mathcal{O}\left((\frac{J'}{T})^3 \right) \\
&= 2 \, (\frac{J'}{T})^2 \, \Big[2 \, \chi_{\text{1d-Ising}}(2n - 2m) + \chi_{\text{1d-Ising}}(2n - 2m + 1) \\
&\qquad\qquad + \chi_{\text{1d-Ising}}(2n - 2m - 1) \Big] + \mathcal{O}\left((\frac{J'}{T})^3 \right)
\end{aligned} \qquad \text{(D.2)}$$

Here $\chi_{\text{1d-Ising}}(n)$ is the correlation function of the one-dimensional Ising chain,

$$\chi_{\text{1d-Ising}}(n) = (-)^n \left(\tanh(\frac{J}{T}) \right)^{|n|} \; . \qquad \text{(D.3)}$$

Hence

$$\chi(2m+1, 2n+1) = -\frac{8(J'/T)^2}{e^{4J/T} - 1} \left(\tanh(\frac{J}{T}) \right)^{2|n-m|} + \mathcal{O}\left((\frac{J'}{T})^3 \right) \; . \qquad \text{(D.4)}$$

In the limit considered here, the correlation between any pair of middle spins of the same row is negative for all distances. It is mediated by the interaction with the neighbouring spin chains.

References

[1] Yavors'kii T, Apel W and Everts H U 2007 *Phys. Rev. B* **76** 064430

[2] Wang F, Vishwanath A and Kim Y B 2007 *Phys. Rev. B* **76** 094421

[3] Schnyder A P, Starykh O A and Balents L 2008 *Phys. Rev. B* **78** 174420

[4] Kano K and Naya S 1953 *Prog. Theor. Phys.* **10** 158

[5] Li W, Gong S S, Zhao Y, Ran S J, Gao S and Su G 2010 *Phys. Rev. B* **82** 134434

[6] Apel W 1986 *Z. Phys.* **63** 185–188

[7] Montroll E W, Potts R B and Ward J C 1962 *J. Math. Phys.* **4** 308

[8] Stephenson J 1964 *J. Math. Phys.* **5** 1009

[9] Schulz T D, Mattis D C and Lieb E H 1964 *Rev. Mod. Phys.* **36** 856

[10] Wu T T 1966 *Phys. Rev.* **149** 380

[11] Stephenson J 1970 *J. Math. Phys.* **11** 413,420

[12] Au-Yang H and McCoy B 1974 *Phys. Rev. B* **10** 3885

[13] Gabay M 1980 *J. Physique - LETTRES* **41** L427

[14] Wolf W F and Zittartz J 1982 *Z. Phys. B* **47** 341

[15] Wolf W F and Zittartz J 1982 *Z. Phys. B* **49** 229

[16] McCoy B M and Wu T T 1973 *The two-dimensional Ising model* (Cambridge, Massachusetts, USA: Harvard University Press)

[17] Grenander U and Szegö G 1958 *Toeplitz Forms and their Applications* (Berkely and Los Angeles: University of California Press)

[18] Geilikman M B 1974 *Sov. Phys. JETP* **39** 570

[19] Vdovichenko N V 1965 *Sov. Phys. JETP* **20** 477

[20] Vdovichenko N V 1965 *Sov. Phys. JETP* **21** 350

[21] Jordan P and Wigner E 1928 *Z. Physik* **47** 631

[22] Berezin F A 1966 *The Method of Second Quantization* (New York: Acad. Press) ISBN 0120894505

[23] Rraján P 1985 (*Lecture Notes in Physics* vol 226) (Heidelberg: Springer) chap Exact disorder solutions, p 286